

Acne phototherapy with a high-intensity, enhanced, narrowband, blue light source: an open study and in vitro investigation

Akira Kawada *, Yoshinori Aragane, Hiroko Kameyama, Yoshiko Sangen, Tadashi Tezuka

Department of Dermatology, Kinki University School of Medicine, Ohno-Higashi 377-2, Osaka-Sayama city, Osaka 589-8511, Japan

Received 19 April 2002; received in revised form 17 June 2002; accepted 19 June 2002

Spectroscopic Study of Porphyrin Production and Accumulation in Culture of Propionibacterium acnes

M. Idrish Miah, Department of Physics, Bangladesh Institute of Technology, Chittagong, Chittagong-4349, Bangladesh

Biotechnology, Volume 1 Number 1: 21-27, 2002

© Asian Network for Scientific Information

Phototherapy in the Treatment of Acne Vulgaris. What is its Role?

Aikaterini Charakida,¹ Edward D. Seaton,¹ Marietta Charakida,² Paul Mouser,¹ Antonias Avgerinos³ and Antony C. Chu¹

¹ Department of Dermatology, Faculty of Medicine, Hammersmith Campus, Imperial College, London, UK

² Vascular Physiology Unit, Great Ormond Street Hospital, Institute of Child Health, London, UK

³ Elena Venizelou Hospital, Athens, Greece

Am J Clin Dermatol 2004; 5 (4): 211-216

An open study to determine the efficacy of blue light in the treatment of mild to moderate acne: preliminary data

C.A. MORTON¹, R.D. SCHOLEFIELD², C. WHITEHURST³, J. BIRCH³

¹. Dermatology dept, Falkirk Royal Infirmary

². Sequani Consumer Ltd, Vine House, West New Street, Ledbury, Herefordshire

³. Photo Therapeutics Ltd, Station House, Stamford New Road, Altrincham

In-vivo-Porphyrinproduktion von Propionibacterium acnes unter systemischer Aknetherapie mit Isotretinoin und Minozyklin

Klinik und Poliklinik für Dermatologie und Allergologie der Ludwig-Maximilians-Universität München, Direktor: Prof. Dr. med. Dr. h.c. G. Plewig

vorgelegt von Kathrin Merk aus Garmisch-Partenkirchen 2004

Visible light promotes proliferation of normal skin cells, Investigation

by Professor Rachel LUBART, Department of Physics, Bar-Ilan University, Tel-Aviv, Israel

Effect of Visible Light on Reactive Oxygen Species Production

R. Lubart,¹ H. Friedmann,¹ R. Lavie,¹ N. Grossman,² M. Sinyakov and S. Belotsky

Department of Chemistry and Physics, Department of Life Sciences

Bar-Ilan University, Ramat-Gan 52900, Israel

*) Alle aufgeführten Studien, Abstracts und Conclusions können auf Wunsch zur Verfügung gestellt werden.

Studien*) zu Indikationen: Akne (II)

An action spectrum for blue and near ultraviolet inactivation of Propionibacterium acnes; with emphasis on a possible porphyrin photosensitisation.

Kjeldstad B, Jhonsson, Photochemistry-Photobiology 1986; 43(1); 67-70

Singlet oxygen (1 delta g) generation from coproporphyrin in Propionibacterium acnes on irradiation.

Arakane K, Ryu A, Hayashi C, Masunaga T, Schinmoto K, Mashiko S, Nagano T, Hirobe M., Biochem Biophys Res Commun 1996; 223 (3): 578-82.

Improvement of rheologic parameters, ligand- and oxygen-binding capacity of erythrocytes of circulating blood after exposure of the body surface to visible polarized light.

K.A.Samoilova, K.D.Obolenskaya, A.V.Vologdina, N.V.Mineeva, N.Yu.Romanenko, M.F.Balljuzek: Published: 8th Congress of European Society for Photobiology. Book of Abstracts, P106, p.145, Granada (1999)

Phototherapy with blue and red light in the treatment of acne vulgaris

Papageorgiou P, Katsambas A, Chu A.
Unit of Dermatology, Imperial College of Science, Technology and Medicine,
Hammersmith Hospital, DuCane Road, London W12 0NN, U.K.
2000 May; 142(5):973-8. Related Articles

Wissenschaftliche, fachdermatologische, gutachterliche Stellungnahme zur Behandlung von Akne vulgaris

Prof. Dr. med.habil. Lutz Kowalzik, HUMAINE Vogtland Klinikum Plauen,07/2005

*) Alle aufgeführten Studien, Abstracts und Conclusions können auf Wunsch zur Verfügung gestellt werden.